


Supplementary Materials for

**Minfi: A flexible and comprehensive Bioconductor package for the analysis of Infinium DNA Methylation microarrays**

Martin J. Aryee, Andrew E. Jaffe, Hector Corrada-Bravo, Christine Ladd-Acosta, Andrew P. Feinberg, Kasper D. Hansen\*, Rafael A. Irizarry\*

Corresponding authors:


\* Rafael A. Irizarry: [rafa@jimmy.harvard.edu](mailto:rafa@jimmy.harvard.edu); \* Kasper D. Hansen: [khansen@jhsph.edu](mailto:khansen@jhsph.edu)




**Supplementary Figure 1.** As Figure 3A but for placenta replicates: for each locus we compute the average and standard deviation (SD) across placenta technical samples. We fit a loess curve to the SD versus average scatterplot for each method. The resulting curve is shown for each preprocessing method.

# Liver

# Placenta




**Supplementary Figure 2.** As Supplementary Figure 1 but for both tissues and stratified by probe type and region: for each locus we compute the average and standard deviation (SD) across technical samples. We fit a loess curve to the SD versus average scatterplot for each method. The resulting curve is shown for each preprocessing method




**Supplementary Figure 3.** As Supplementary Figure 1 but for the results of applying the algorithms to the data on the logit scale: for each locus we compute the average and standard deviation (SD) across technical samples. We fit a loess curve to the SD versus average scatterplot for each method. The resulting curve is shown for each preprocessing method.

# Liver



# Placenta



**Supplementary Figure 4.** As Supplementary Figure 3 but stratified by probe type and region: for each locus we compute the average and standard deviation (SD) across technical samples. We fit a loess curve to the SD versus average scatterplot for each method. The resulting curve is shown for each preprocessing method.



**Supplementary Figure 5.** Distributions of beta methylation values stratified by probe type (Type I and II), genomic region, and preprocessing method.



**Supplementary Figure 6.** Genotype effects on DNA methylation levels. A) We computed the differences in average methylation between Caucasian and Hispanic ethnicities and plot a quantile-quantile plot of the SNP-affected probes versus the non-affected probes. B) Each SNP-affected probe is paired with its closest non-affected probe. We plot the Caucasian-Hispanic differences of the SNP affected probes against the difference at its closest non-affected probe.